Anisotropic Surface Meshing

Siu-Wing Cheng*

Tamal K. Dey†

Edgar A. Ramos‡

Rephael Wenger§

Abstract

We study the problem of triangulating a smooth closed implicit surface Σ endowed with a 2D metric tensor that varies over Σ . This is commonly known as the anisotropic surface meshing problem. We extend the 2D metric tensor naturally to 3D and employ the 3D anisotropic Voronoi diagram of a set P of samples on Σ to triangulate Σ . We prove that a restricted dual, $\text{Mesh } P$, is a valid triangulation homeomorphic to Σ under appropriate conditions. We also develop an algorithm for constructing P and $\text{Mesh } P$. In addition to being homeomorphic to Σ , each triangle in $\text{Mesh } P$ is well-shaped when measured using the 3D metric tensors of its vertices. Users can set upper bounds on the anisotropic edge lengths and the angles between the surface normals at vertices and the normals of incident triangles (measured both isotropically and anisotropically).

1 Introduction

Many applications in science and engineering need to mesh a smooth closed surface Σ for numerical simulations, prototyping, rendering and various other purposes. A variety of algorithms and software systems based on them have been developed for the problem by the mesh generation, computer graphics, and image processing communities, e.g. [4, 12, 15, 17, 19, 21]. A subset of them focus on implicit surfaces, e.g. [4, 15, 17, 19]. An implicit surface representation is appealing for the relative ease in combining surfaces, computing line-surface intersection, and carrying out the inside/outside test. In this representation, there is an implicit function $E: \mathbb{R}^3 \rightarrow \mathbb{R}$ and Σ is its zero-set $E(x) = 0$. The surface Σ can have arbitrary genus.

Despite the experimental success, it is not until recently that some algorithms have been designed with a guarantee about output quality: specifically, the out-

put triangulation being homeomorphic to the input surface Σ and being geometrically close to it. Cheng, Dey, Edelsbrunner, and Sullivan [10] designed a meshing algorithm for skin surfaces in molecular modeling. Boissonnat and Oudot [6] and Cheng, Dey, Ramos, and Ray [11] proposed algorithms for general implicit surfaces. Although both algorithms sample points and compute a restricted Delaunay triangulation, they differ in many ways. In particular, a lower bound estimation of the local feature size is needed in [6] where critical point computations replacing local feature size are required in [11]. Boissonnat, Cohen-Steiner, and Vetter [7] designed an algorithm to approximate the level sets of the implicit function, which yields an isotopic surface triangulation.

The above certified meshing algorithms are designed for the isotropic setting in which nearly equilateral triangles are used. Sometimes, long and skinny triangles offer better approximation because they adapt to the principal curvatures of the surface. They are also preferred in numerical simulations where the physical phenomena are strongly directional. These applications call for an *anisotropic triangulation*. The general formulation is to associate a 2D *metric tensor* M_x , a 2×2 positive definite symmetric matrix, with each point $x \in \Sigma$. Its eigenvalues $0 < k_{x,1} \leq k_{x,2}$ and corresponding orthogonal unit eigenvectors $d_{x,1}$ and $d_{x,2}$ specify the following deformation on the tangent plane at x . Any point z on the tangent plane is mapped to the point $x + \langle z - x, d_{x,1} \rangle \cdot \sqrt{k_{x,1}} d_{x,1} + \langle z - x, d_{x,2} \rangle \cdot \sqrt{k_{x,2}} d_{x,2}$. So the distance components along $d_{x,1}$ and $d_{x,2}$ are scaled by factors $\sqrt{k_{x,1}}$ and $\sqrt{k_{x,2}}$, respectively. We call $\sqrt{k_{x,1}}$ and $\sqrt{k_{x,2}}$ the *scaling factors* and $d_{x,1}$ and $d_{x,2}$ the *principal directions*. The points on the tangent plane at distance 1 from x (measured using M_x) lie on an ellipse centered at x with $d_{x,1}$ as the major axis. To adapt to M_x , the surface mesh triangles near x should be elongated in direction $d_{x,1}$.

Few theoretical results are known for the anisotropic triangulation problem. Algorithms were obtained recently for the problem of computing a 2D conformal anisotropic triangulation of a planar straight line graph with no angle much less $\pi/2$ [8, 16]. The definition of anisotropic Voronoi diagram originates from the work of Labelle and Shewchuk [16]. A surface associated with

*Supported by the Research Grant Council, Hong Kong, China (HKUST6181/04E). Department of Computer Science, HKUST, Hong Kong. Email: scheng@cs.ust.hk.

†Supported by NSF CCR-0430735 and NSF DMS-0310642. Department of Computer Science and Engineering, Ohio State University, USA. Email: tamaldey@cse.ohio-state.edu.

‡Department of Computer Science, University of Illinois at Urbana-Champaign, USA. Email: eramosn@cs.uiuc.edu.

§Department of Computer Science and Engineering, Ohio State University, USA. Email: wenger@cse.ohio-state.edu.

a varying metric tensor is a Riemannian manifold. Leiben and Letscher [18] showed that if the sampling density is sufficiently high, the dual of the Voronoi diagram defined using the shortest path distance is a valid triangulation. Unfortunately, this distance calculation is computationally difficult and it is unclear how to obtain the samples in the first place.

In this paper, we assume that the largest scaling factor at any point $x \in \Sigma$ is bounded from above, but we do not need to know the bound. We assume an oracle to return M_x for each point $x \in \Sigma$, but we do not need a closed form formula for generating M_x . The metric tensor should vary smoothly and we will make this concrete later. We extend M_x to a 3D metric tensor by adding a distance component with scaling factor $\sqrt{k_{x,2}}$ in the normal direction at x . On planes parallel to the tangent plane at x , the 3D metric tensor measures angles and lengths identically as M_x . Then we employ the 3D anisotropic Voronoi diagram of a set P of samples on Σ to triangulate Σ . It is known that the dual of a 3D anisotropic Voronoi diagram may not be a valid tetrahedralization [8, 16]. However, we show that a restricted dual, $\text{Mesh } P$, is a valid triangulation homeomorphic to Σ under appropriate conditions. (See Theorem 5.1.)

We develop an incremental point insertion algorithm to grow P as in the isotropic case [11]. The final $\text{Mesh } P$ is homeomorphic to Σ . Users can specify two upper bounds ℓ and ζ such that the edge length is at most ℓ when measured using the 3D metric tensors of the edge endpoints, and the surface normal at a vertex makes an angle at most ζ with the normal of any incident triangle. The bound ζ also holds when the angle is measured using the 3D metric tensor of the vertex. Each triangle in $\text{Mesh } P$ has bounded aspect ratio when measured using the 3D metric tensors of its vertices. When ζ is small, a triangle will be almost parallel to the tangent planes at its vertices. Then bounded aspect ratio in the anisotropic setting implies that for each triangle vertex x , the triangle is elongated more or less in the direction $d_{x,1}$ as desired.

2 Metric extension and preliminaries

Vector and matrix. We use $\|\cdot\|$ to denote the l^2 -norm and the matrix 2-norm. So $\|(a, b, c)^t\| = \sqrt{a^2 + b^2 + c^2}$. For a real symmetric matrix A , take the absolute values of its eigenvalues and let $\lambda_{\max}$ and $\lambda_{\min}$ be the largest and smallest ones, respectively. Then $\|A\| = \lambda_{\max}$ and $\|A^{-1}\| = 1/\lambda_{\min}$.

3D metric tensor. At each point $x \in \Sigma$, we define a 3D metric tensor $K_x = U_x^t \cdot D_x \cdot U_x$, where D_x is a 3×3 diagonal matrix with entries $k_{x,1}$, $k_{x,2}$, and $k_{x,3}$ from top

left to bottom right. The matrix U_x is a 3×3 matrix and it transforms space so that the principal directions and the unit outward normal at x become the coordinate system. The new entry $k_{x,3}$ should be chosen so that $k_{x,3} \geq k_{x,2}$. A natural choice is $k_{x,3} = k_{x,2}$ and this corresponds to inflating the “unit ellipse” on the tangent plane at x into a “unit ellipsoid” in 3D. If desired, one can also choose $k_{x,3}$ to be greater than $k_{x,2}$. Our results hold as long as $\max_{x \in \Sigma} k_{x,3}$ is bounded from above. One can express $K_x = F_x^t \cdot F_x$, where $F_x = U_x^t \cdot D'_x \cdot U_x$ and the diagonal entries of D'_x are $\sqrt{k_{x,1}}$, $\sqrt{k_{x,2}}$, and $\sqrt{k_{x,3}}$ from top left to bottom right. From the perspective of x , the distance between two points $a, b \in \mathbb{R}^3$ is $\|F_x \cdot (b - a)\|$.

3D Linear map. For any point $x \in \Sigma$, we define a map φ_x such that for any point $b \in \mathbb{R}^3$, $\varphi_x(b) = F_x \cdot b$. For any subset $B \subseteq \mathbb{R}^3$, we use $\varphi_x(B)$ to denote $\{\varphi_x(b) : b \in B\}$. We call $\mathbb{R}^3$ the *primal space* and $\varphi_x(\mathbb{R}^3)$ the φ_x -*space*. We extend the notation $\varphi_x(\cdot)$ in several ways to ease the presentation. If we write a vector $d \in \mathbb{R}^3$ as a triple of real numbers, then $F_x \cdot d$ is the vector in the φ_x -space that d is mapped to. We use $\varphi_x(d)$ to denote $F_x \cdot d$. Given two vectors d_1 and d_2 , we use $\angle d_1, d_2$ to denote the angle between d_1 and d_2 and we use $\varphi_x(\angle d_1, d_2)$ to denote $\angle \varphi_x(d_1), \varphi_x(d_2)$. Given three points a, b, c , we use $\angle abc$ to denote $\angle a - b, c - b$. Given two points $a, b \in \mathbb{R}^3$, we use $d_x(a, b)$ to denote the distance $\|\varphi_x(a) - \varphi_x(b)\|$ in the φ_x -space. In the φ_x -space, x 's distance metric becomes isotropic. If y 's distance metric is nearly isotropic in the φ_x -space, the bisector surface of $\varphi_x(x)$ and $\varphi_x(y)$ intersects the segment $\varphi_x(xy)$ at nearly right angle and the bisector surface is nearly perpendicular to $\varphi_x(xy)$ near the intersection point. Collinearity and incidence are preserved by φ_x . If a plane H is tangent to Σ , $\varphi_x(H)$ is also tangent to $\varphi_x(\Sigma)$. Angles are not preserved in general. But if d_1 is a normal vector at x and d_2 is orthogonal to d_1 , then $\varphi_x(d_1)$ and $\varphi_x(d_2)$ are also orthogonal.

Propagating the metric. The *medial axis* $\mathcal{X}$ of Σ is the set of centers of maximal balls empty of points on Σ . For any point $x \in \Sigma$, the *local feature size* $f(x)$ is the distance from x to $\mathcal{X}$. For any point $p \in \mathbb{R}^3 \setminus \mathcal{X}$, we use $\tilde{p}$ to denote the closet point on Σ to p . Define Σ^* to be the set of points p whose distance from Σ is at most $f(\tilde{p})/2$. For each $p \in \Sigma^*$, define $F_p = F_{\tilde{p}}$ and $K_p = K_{\tilde{p}}$. The set of points $\Sigma^* \setminus \Sigma$ and their matrices are only needed for analysis and our algorithm never computes them.

Distortion. The *relative distortion* between two points $p, q \in \Sigma^*$ is $\tau_{pq} = \max\{\|F_p \cdot F_q^{-1}\|, \|F_q \cdot F_p^{-1}\|\}$. We have $d_p(a, b) = \|F_p \cdot (a - b)\| \leq \|F_p \cdot F_q^{-1}\| \cdot d_q(a, b) \leq \tau_{pq} d_q(a, b)$. Conversely, we also have $d_p(a, b) \geq$

$d_q(a, b)/\tau_{pq}$. Define $\tau_* = (65/64)^{1/4}$. For each point $p \in \Sigma^*$, let $\gamma(p)$ denote $\max\{\lambda \in \mathbb{R} : \forall q \in \Sigma^*, d_p(p, q) \leq \lambda \Rightarrow \tau_{pq} \leq \tau_*\}$. We assume that F_x varies smoothly over $x \in \Sigma$ so that $\min_{p \in \Sigma^*} \gamma(p)$ is strictly positive. For any points $x, y \in \Sigma$, we define $f_x(y)$ to be the distance from $\varphi_x(y)$ to the medial axis of $\varphi_x(\Sigma)$ in the φ_x -space. Define $\gamma = \min\{\min_{p \in \Sigma^*} \gamma(p), \min_{x \in \Sigma} f_x(x)\}$. Thus if $d_x(x, y) \leq \gamma$, then $\tau_{xy} \leq \tau_*$. The value γ defines a neighborhood of x in which the geometry and the metric tensor do not vary much. The value γ is only needed for analysis and our algorithm never computes it.

Anisotropic Voronoi diagram. Let P be a finite set of points on Σ . The *anisotropic Voronoi cell* of $u \in P$ is $V_u(P) = \{p \in \mathbb{R}^3 : \forall v \in P, d_u(p, u) \leq d_v(p, v)\}$. For $0 \leq j \leq 3$, the closed faces shared by $4 - j$ Voronoi cells are called *j-dimensional Voronoi faces*. We denote a 2-dimensional *Voronoi facet* $V_u(P) \cap V_v(P)$ by $V_{uv}(P)$ and a 1-dimensional *Voronoi edge* $V_u(P) \cap V_v(P) \cap V_w(P)$ by $V_{uvw}(P)$. The *anisotropic Voronoi diagram* $\text{Vor } P$ is the collection of Voronoi cells together with their boundary faces. The diagram $\text{Vor } P$ is the lower envelope of $|P|$ surfaces in $\mathbb{R}^4$. It has $O(|P|^{4+\epsilon})$ complexity [20]. The *restricted dual*, $\text{Mesh } P$, is a simplicial complex comprising of the points in P , the edges uv such that $V_{uv}(P) \cap \Sigma \neq \emptyset$, and the triangles uvw such that $V_{uvw}(P) \cap \Sigma \neq \emptyset$.

Wedge and visibility. Let Q be a set of points on Σ . Define the *wedge* of Q to be $\text{wedge}(Q) = \{x \in \mathbb{R}^3 : \varphi_{v_i}(\angle x v_i v_j) < \pi/2, \forall v_i, v_j \in Q\}$. By definition, $\text{wedge}(Q)$ is an open set.

LEMMA 2.1. *Let $\tau > 1$ be a constant. Let v_1 and v_2 be points on Σ . Let x be a point in $V_{v_1 v_2}(\{v_1, v_2\})$.*

- (i) *If $x \in \text{wedge}(\{v_1, v_2\})$, the interior of the segment $v_i x$ lies strictly inside $V_{v_i}(\{v_1, v_2\})$.*
- (ii) *If $x \notin \text{wedge}(\{v_1, v_2\})$ and $\tau_{v_1 v_2} \leq \tau$, then $d_{v_i}(v_i, x) \geq \frac{1}{\sqrt{\tau^2 - 1}} \cdot \min\{d_{v_1}(v_1, v_2), d_{v_2}(v_1, v_2)\}$ for $1 \leq i \leq 2$.*
- (iii) *Let Q be a set of points on Σ such that $\tau_{vw} \leq \tau$ for any $v, w \in Q$. If the segment vp does not lie inside $V_v(Q)$ for some point p on the boundary of $V_v(Q)$, then $d_v(v, p) \geq \frac{1}{\sqrt{\tau^2 - 1}} \cdot \min\{d_v(v, w), d_w(v, w)\}$ for some $w \in Q$.*

Proof. (i) and (ii) have been proved in [16]. Consider (iii). As we walk from v towards p along vp , we must leave $V_v(Q)$ and then reenter $V_v(Q)$ at some point q . The point q lies on $V_{vw}(Q)$ for some $w \in Q$, and vq does not lie inside $V_v(\{v, w\})$. By (i), $q \notin \text{wedge}(\{v, w\})$. Then by (ii), $d_v(v, p) \geq d_v(v, q) \geq$

$$\frac{1}{\sqrt{\tau^2 - 1}} \cdot \min\{d_v(v, w), d_w(v, w)\}. \quad \square$$

3 Surface normal variation

For any point $v, x \in \Sigma$, we use $\mathbf{n}(x)$ to denote the unit outward normal of Σ at x and $\mathbf{n}_v(x)$ to denote the unit outward normal of $\varphi_v(\Sigma)$ at $\varphi_v(x)$. Note that $\mathbf{n}_x(x) = \mathbf{n}(x)$ for any point $x \in \Sigma$. The main results in this section, as stated in Lemma 3.2 below, are concerned with the variation of the surface normal. Lemma 3.2(i) states that if two points $v_1, v_2 \in \Sigma$ are very near in the φ_{v_1} -space, the segment $\varphi_{v_1}(v_1 v_2)$ is nearly orthogonal to $\mathbf{n}_{v_1}(v_1)$. The first half of Lemma 3.2(ii) states that if v_1 and v_2 are very near in the φ_v -space, then $\angle \mathbf{n}_v(v_1), \mathbf{n}_v(v_2)$ is small. These results are the anisotropic versions of known results in the isotropic setting [2, 10]. Their proofs are the same too. Lemma 3.2(ii) states further that if v_1 and v_2 are very near in the φ_{v_1} -space, then $\angle \mathbf{n}(v_1), \mathbf{n}(v_2)$ is small. This requires transferring the normal deviation in the φ_{v_1} -space to the primal space, although v_1 and v_2 may be far apart in the primal space. The following result enables the transfer.

LEMMA 3.1. *Let x be a point on Σ . Let H be a plane in the primal space. Let d_1 denote a vector normal to H that makes an acute angle with $\mathbf{n}(x)$. Let d_2 denote a vector normal to $\varphi_x(H)$ that makes an acute angle with $\mathbf{n}_x(x)$. Then $\angle \mathbf{n}(x), d_1 \leq \angle \mathbf{n}_x(x), d_2$.*

Proof. Orient $\mathbb{R}^3$ so that the principal directions and the outward surface normal at x form the coordinate system. Then $\mathbf{n}_x(x) = \mathbf{n}(x) = (0, 0, 1)^t$ points vertically upward. By our orientation of $\mathbb{R}^3$, F_x is a diagonal matrix and its diagonal entries are $\sqrt{k_{x,1}}, \sqrt{k_{x,2}}$ and $\sqrt{k_{x,3}}$ from top left to bottom right. Assume that $d_2 = (a, b, c)^t$. For any vector $u = (e_1, e_2, e_3)^t$ parallel to the plane H , $\langle d_2, F_x \cdot u \rangle = \sqrt{k_{x,1}} a e_1 + \sqrt{k_{x,2}} b e_2 + \sqrt{k_{x,3}} c e_3 = \langle F_x \cdot d_2, u \rangle$. Observe that $F_x \cdot u$ is a vector parallel to $\varphi_x(H)$. Thus, $\langle d_2, F_x \cdot u \rangle = \langle F_x \cdot d_2, u \rangle = 0$. Since this holds for any vector u parallel to H , $F_x \cdot d_2$ is normal to H . Consider $\cos \angle \mathbf{n}_x(x), d_2 = \langle \mathbf{n}_x(x), d_2 \rangle / (\|\mathbf{n}_x(x)\| \cdot \|d_2\|) = c / \sqrt{a^2 + b^2 + c^2}$. Since $\angle \mathbf{n}_x(x), d_2 < \pi/2$ by assumption, we conclude that $c > 0$. If $F_x \cdot d_2$ and d_1 have opposite orientations, then $\cos \angle \mathbf{n}(x), d_1 = \langle \mathbf{n}(x), -(F_x \cdot d_2) \rangle / (\|\mathbf{n}(x)\| \cdot \|F_x \cdot d_2\|) = -c \sqrt{k_{x,3}} / D$, where $D = \sqrt{k_{x,1} a^2 + k_{x,2} b^2 + k_{x,3} c^2}$. But then $\cos \angle \mathbf{n}(x), d_1$ is negative, contradicting the assumption that $\angle \mathbf{n}(x), d_1 \leq \pi/2$. So $\cos \angle \mathbf{n}(x), d_1 = c \sqrt{k_{x,3}} / D \geq c / \sqrt{a^2 + b^2 + c^2} = \cos \angle \mathbf{n}_x(x), d_2$. Hence, $\angle \mathbf{n}(x), d_1 \leq \angle \mathbf{n}_x(x), d_2$. $\square$

Recall that for any point on Σ , the value γ defines a neighborhood in which the geometry and the metric tensor do not vary much.

LEMMA 3.2. *Define the function $\alpha(c) = c/(1-4c)$. Let v_1, v_2 be points in Σ .*

- (i) *Let $\phi = \angle \varphi_{v_1}(v_2 - v_1), \mathbf{n}_{v_1}(v_1)$. Then $d_{v_1}(v_1, v_2) \geq 2\gamma \cos(\min\{\phi, \pi - \phi\})$.*
- (ii) *Let v be a point in Σ . If $d_v(v_1, v_2) \leq c\gamma$ for some constant $c < 1/4$, then $\angle \mathbf{n}_v(v_1), \mathbf{n}_v(v_2) \leq \alpha(c)$. If $v = v_1$, then $\angle \mathbf{n}(v_1), \mathbf{n}(v_2) \leq \angle \mathbf{n}_{v_1}(v_1), \mathbf{n}_{v_1}(v_2) \leq \alpha(c)$.*

Proof. Applying the original proof in [10] in the φ_{v_1} -space proves (i). Consider (ii). Applying the original proof in [2] in the φ_v -space bounds $\angle \mathbf{n}_v(v_1), \mathbf{n}_v(v_2)$. Suppose that $v = v_1$. Let B be the ball in the φ_{v_1} -space centered at $\varphi_{v_1}(v_1)$ with radius $c\gamma$. Because the radius of B is less than $f_{v_1}(v_1)$, it is known that $B \cap \varphi_{v_1}(\Sigma)$ is a topological disk [5]. We go back to the primal space. Let v_2 be a point in $\varphi_{v_1}^{-1}(B) \cap \Sigma$. We first show that $\angle \mathbf{n}(v_1), \mathbf{n}(v_2) < \pi/2$. Assume to the contrary that $\angle \mathbf{n}(v_1), \mathbf{n}(v_2) \geq \pi/2$. Since $\varphi_{v_1}^{-1}(B) \cap \Sigma$ is connected, there is a path from v_1 to v_2 in $\varphi_{v_1}^{-1}(B) \cap \Sigma$. So there is a point $v_3 \in \varphi_{v_1}^{-1}(B) \cap \Sigma$ on this path such that $\angle \mathbf{n}(v_1), \mathbf{n}(v_3) = \pi/2$. But then $\mathbf{n}_{v_1}(v_3)$ is also orthogonal to $\mathbf{n}_{v_1}(v_1)$, contradicting the fact that $\angle \mathbf{n}_{v_1}(v_1), \mathbf{n}_{v_1}(v_3) \leq \alpha(c)$ as $d_{v_1}(v_1, v_3) \leq c\gamma$. So we have $\angle \mathbf{n}(v_1), \mathbf{n}(v_2) < \pi/2$ and $\angle \mathbf{n}_{v_1}(v_1), \mathbf{n}_{v_1}(v_2) \leq \alpha(c) < \pi/2$. By applying Lemma 3.1 to v_1 and the tangent plane at v_2 , we get $\angle \mathbf{n}(v_1), \mathbf{n}(v_2) \leq \angle \mathbf{n}_{v_1}(v_1), \mathbf{n}_{v_1}(v_2)$. $\square$

4 Triangles

Let $t = v_1 v_2 v_3$ be a triangle with vertices on Σ . We use $\mathbf{n}(t)$ to denote the normal of t in the primal space that makes an acute angle with $\mathbf{n}(v_i)$, where v_i is the vertex with the largest angle in t . Similarly, for any point $v \in \Sigma$, $\mathbf{n}_v(t)$ denotes the normal of $\varphi_v(t)$ that makes an acute angle with $\mathbf{n}_v(v_j)$, where $\varphi_v(v_j)$ is the vertex with the largest angle in $\varphi_v(t)$. We use $\text{length}(t)$ to denote $\min_{1 \leq i \leq 3} s_i$, where s_i is the minimum side length of $\varphi_{v_i}(t)$. The main results in this section are that under suitable conditions, $\varphi_{v_i}(t)$ is well-shaped (Lemma 4.1), both $\angle \mathbf{n}_{v_i}(t), \mathbf{n}_{v_i}(v_i)$ and $\angle \mathbf{n}(t), \mathbf{n}(v_i)$ are small (Lemma 4.2), and the dihedral angles between adjacent triangles are close to π (Lemma 4.4). The analogous version of Lemma 4.1 in the plane has been proved in [16].

In the isotropic setting, a triangle is well-shaped if its *radius-edge ratio* (the ratio of circumradius to

shortest edge length) is bounded. In our case, the ratio $d_{v_i}(v_i, z)/\text{length}(t)$, where $z \in V_t(\{v_1, v_2, v_3\})$, plays a critical role. Recall that τ_* denotes the constant $(65/64)^{1/4}$.

LEMMA 4.1. *Let $\rho \in (1, \sqrt{2})$ be a constant. Let $t = v_1 v_2 v_3$ be a triangle with vertices on Σ . Assume that $d_{v_i}(v_i, z) \leq 4\rho \cdot \text{length}(t)$ for some point $z \in V_t(\{v_1, v_2, v_3\})$. Let $v \in \Sigma$ be a point such that $\tau_{vv_i} \leq \tau_*$ for $1 \leq i \leq 3$. Then for each angle ϕ of t , $\varphi_v(\phi) \geq \sin^{-1}(\chi)$, where $\chi = \frac{1-32(\tau_*^4-1)}{128\tau_*^2}$. Hence, the radius-edge ratio of $\varphi_v(t)$ is at most $\rho_1 = 2/\chi$.*

Proof. We go to the φ_v -space. Let $a_i = \varphi_v(v_i)$ for $1 \leq i \leq 3$. Let $y = \varphi_v(z)$. Let x be the projection of y onto the plane of $\varphi_v(t)$.

Each triangle edge incident to a_i bounds an open halfplane disjoint from $\varphi_v(t)$. We call the intersection of these two open halfplanes the negative cone of a_i . We claim that x does not lie in the negative cone of any a_i . Assume to the contrary that x lies in the negative cone of a_1 . Then $\angle a_2 a_1 x$ or $\angle a_3 a_1 x$ is at least $\pi/2$, say $\angle a_2 a_1 x$. So $\angle a_2 a_1 y \geq \pi/2$ too. This implies that $|a_2 y|^2 \geq |a_1 y|^2 + |a_1 a_2|^2$. Since $\tau_{vv_i} \leq \tau_*$ for all v_i , we have $|a_2 y| \leq \tau_* d_{v_2}(v_2, z) = \tau_* d_{v_1}(v_1, z)$, $|a_1 y| \geq d_{v_1}(v_1, z)/\tau_*$, and $|a_1 a_2| \geq d_{v_1}(v_1, v_2)/\tau_*$. It follows that $\tau_*^4 d_{v_1}(v_1, z)^2 \geq d_{v_1}(v_1, z)^2 + d_{v_1}(v_1, v_2)^2$. But then $d_{v_1}(v_1, z) \geq d_{v_1}(v_1, v_2)/\sqrt{\tau_*^4 - 1} = 8d_{v_1}(v_1, v_2) > 4\rho \cdot d_{v_1}(v_1, v_2)$, contradicting our assumptions.

By our claim, there exists a vertex a_i such that $a_i x$ stabs $a_1 a_2 a_3$, say $a_2 x$. Let $r = d_{v_i}(v_i, z)$ for $1 \leq i \leq 3$. The following inequalities will be useful later:

$$\begin{aligned} |a_i y| &\leq \tau_* r. \\ |a_i y| &\geq r/\tau_*. \\ |a_i a_j| &\geq d_{v_i}(v_i, v_j)/\tau_* \geq r/(4\tau_* \rho). \end{aligned}$$

Note that $\tau_* + 1/\tau_* > 1/(4\tau_* \rho)$ and $1/(4\tau_* \rho) + 1/\tau_* > \tau_*$. This implies that one can form a triangle $b_y b_i b_j$ such that $|b_i b_y| = r/\tau_*$, $|b_i b_j| = r/(4\tau_* \rho)$, and $|b_j b_y| = \tau_* r$. Since $b_j b_y$ is the longest edge in the triangle $b_y b_i b_j$, both $\angle b_i b_y b_j$ and $\angle b_i b_j b_y$ are acute. Since $|b_i b_y| \leq |a_i y|$, $|b_i b_j| \leq |a_i a_j|$ and $|b_j b_y| \geq |a_j y|$, by scaling $|b_j b_y|$ down to $|a_j y|$ and considering the circumcircle of $b_y b_i b_j$, one can show that $\angle b_y b_i b_j \geq \angle y a_i a_j$. Applying the law of cosines yields

$$\cos \angle b_y b_i b_j = \frac{1 - 16(\tau_*^4 - 1)\rho^2}{64\tau_*^2 \rho^2} \cdot 8\tau_*^2 \rho.$$

By substituting $\rho \leq \sqrt{2}$, we get $\cos \angle b_y b_i b_j \geq 8\chi\tau_*^2 \rho$. Thus $\angle b_y b_i b_j$ is acute and so is $\angle y a_i a_j$. It follows that $\angle y a_i a_j \leq \cos^{-1}(8\chi\tau_*^2 \rho)$. If $\angle x a_i a_j \geq \pi/2$, then $\angle y a_i a_j$ would be at least $\pi/2$ too, which is impossible. Thus

$\angle x a_i a_j < \pi/2$ and so

$$\angle x a_i a_j \leq \angle y a_i a_j \leq \cos^{-1}(8\chi\tau_*^2\rho).$$

Recall that a_2x stabs $a_1a_2a_3$. The support line of a_2x cuts a_1a_3 at a point c . Without loss of generality, assume that $\angle a_2ca_3 \leq \pi/2$. Then $\angle a_2a_3a_1 \geq \pi/2 - \angle a_2a_3 \geq \pi/2 - \cos^{-1}(8\chi\tau_*^2\rho) = \sin^{-1}(8\chi\tau_*^2\rho)$. In order to lower bound $\angle a_3a_1a_2$ and $\angle a_1a_2a_3$, there are two cases to consider depending on the acuteness of $\angle a_2a_3a_1$. First, for any i and j , $|a_i a_j| \leq d_v(v_i, z) + d_v(v_j, z) \leq \tau_* d_{v_i}(v_i, z) + \tau_* d_{v_j}(v_j, z) = 2\tau_* r$.

Suppose that $\angle a_2a_3a_1 \leq \pi/2$. By the law of sines, $\sin \angle a_3a_1a_2 = (|a_2a_3| \sin \angle a_2a_3a_1) / |a_1a_2|$. Thus $\sin \angle a_3a_1a_2 \geq \frac{r}{4\tau_*\rho} \cdot 8\chi\tau_*^2\rho \cdot \frac{1}{2\tau_*r} = \chi$. Thus, $\angle a_3a_1a_2 \geq \sin^{-1}(\chi)$. Similarly, $\angle a_1a_2a_3 \geq \sin^{-1}(\chi)$.

Suppose that $\angle a_2a_3a_1 > \pi/2$. In this case, x must lie inside the triangle $a_1a_2a_3$. Otherwise, $\angle a_2a_3a_1 \leq \angle x a_3 a_2$ which has been proved to be acute, a contradiction. Since x lies inside the triangle $a_1a_2a_3$, $\angle a_2a_3a_1 \leq \angle x a_3 a_2 + \angle x a_3 a_1 \leq 2\cos^{-1}(8\chi\tau_*^2\rho)$. Since $\angle a_2a_3a_1 > \pi/2$, $\sin \angle a_2a_3a_1$ is minimized when $\angle a_2a_3a_1$ is maximized. Thus, by the law of sines, $\sin \angle a_3a_1a_2 \geq \frac{r}{4\tau_*\rho} \cdot \sin(2\cos^{-1}(8\chi\tau_*^2\rho)) \cdot \frac{1}{2\tau_*r} = 2\chi \sin(\cos^{-1}(8\chi\tau_*^2\rho))$. By our choices of τ_* , χ and ρ , $\sin(\cos^{-1}(8\chi\tau_*^2\rho)) > 1/2$. So $\sin \angle a_3a_1a_2 \geq \chi$, implying that $\angle a_3a_1a_2 \geq \sin^{-1}(\chi)$. Similarly, $\angle a_1a_2a_3 \geq \sin^{-1}(\chi)$. $\square$

From now on, we reserve ρ to express the bound on $d_{v_i}(v_i, z)/\text{length}(t)$, $\sin^{-1}(\chi)$ to express the lower bound on the angles of $\varphi_v(t)$, and $\rho_1 = 2/\chi$ to express the bound on the radius-edge ratio of $\varphi_v(t)$, where v , v_i and t are as described in Lemma 4.1.

Lemma 4.2 below analyzes the normal of t and distance of t to Σ . The original proof in [3] proves (i): bounding $\angle \mathbf{n}_v(t), \mathbf{n}_v(v_i)$ for any point $v \in \Sigma$ nearby. Then we use Lemma 3.1 to prove (ii): transferring this angle bound to the primal space. Lastly, every point in t is close to Σ as stated in (iii).

LEMMA 4.2. Define the function $\beta(c) = \sin^{-1}(c) + \sin^{-1}(2\sin(2\sin^{-1}(c))/\sqrt{3})$. Let $\rho \in (1, \sqrt{2}]$ and $\rho_1 = 2/\chi$ be two constants. Let $t = v_1v_2v_3$ be a triangle with vertices on Σ . Assume that $d_{v_i}(v_i, z) \leq 4\rho \cdot \text{length}(t)$ for some point $z \in V_t(\{v_1, v_2, v_3\})$. For sufficiently small constant c , the following hold:

- (i) if v is a point on Σ with $d_v(v, v_i) \leq c\gamma$ for $1 \leq i \leq 3$, then for $1 \leq i \leq 3$, $\angle \mathbf{n}_v(t), \mathbf{n}_v(v_i) \leq \beta(2\rho_1 c) + \alpha(2c)$;
- (ii) if $d_{v_i}(v_i, v_j) \leq c\gamma$ for $1 \leq i, j \leq 3$, then for $1 \leq i \leq 3$, $\angle \mathbf{n}(t), \mathbf{n}(v_i) \leq \beta(2\rho_1 c) + \alpha(2c)$;

- (iii) if v is a point on Σ with $d_v(v, v_i) \leq c\gamma$ for $1 \leq i \leq 3$, then for any point $p \in t$, $\varphi_v(p) + \lambda \cdot \mathbf{n}_v(v) \in \varphi_v(\Sigma)$ and $p + \lambda' \cdot \mathbf{n}(v) \in \Sigma$ for some λ, λ' such that $|\lambda| < 40\rho_1^2 c^2 \gamma$ and $|\lambda'| < 40\rho_1^2 c\gamma$.

To lower bound the dihedral angle between two triangles $t_1 = v_1v_3v_4$ and $t_2 = v_2v_3v_4$ sharing the edge v_3v_4 , our approach is to prove that the dihedral angle between $\varphi_{v_3}(t_1)$ and $\varphi_{v_3}(t_2)$ is large in the φ_{v_3} -space. Then we transfer this bound to the primal space using Lemma 4.2(ii). We can do this only when t_1 and t_2 are sufficiently small. So we introduce some parameters to control the size of triangles. Let $\omega \in (0, \sqrt{1 - 1/\tau_*})$ be a parameter. Define the following:

- $\tau_\omega = 1/(1 - \omega^2)$,
- $\forall x \in \Sigma^*, c_{\omega, x} = \max\{\lambda \leq 1/2 : \forall y \in \Sigma^*, d_x(x, y) \leq 3\lambda\gamma \Rightarrow \tau_{xy} \leq \tau_\omega\}$,
- $c_\omega = \min\{\omega^3, \min_{x \in \Sigma^*} c_{\omega, x}\}$.

The parameter τ_ω is less than τ_* and τ_ω approaches 1 as ω decreases. This reflects a tighter allowance for metric distortion. Then $c_{\omega, x}\gamma$ measures the neighborhood size of a point $x \in \Sigma^*$ in which the metric distortion is at most τ_ω . Finally, we will use the global minimum $c_\omega\gamma$ to control the triangle size. As ω decreases, $c_\omega\gamma$ approaches zero. In defining c_ω , we take a minimization with ω^3 and this will be useful in the proof of Lemma 5.1.

The dihedral angle bound between $\varphi_{v_3}(t_1)$ and $\varphi_{v_3}(t_2)$ is obtained by showing that the tangents to the dual Voronoi edges of $\varphi_{v_3}(t_1)$ and $\varphi_{v_3}(t_2)$ are nearly orthogonal to $\varphi_{v_3}(\Sigma)$ around $\varphi_{v_3}(v_3)$, and the two dual Voronoi edges intersect the planes of $\varphi_{v_3}(t_1)$ and $\varphi_{v_3}(t_2)$. The exact statements are given in Lemma 4.3 below. Its proof is omitted.

LEMMA 4.3. Let $\rho = (1, \sqrt{2}]$ and $\rho_1 = 2/\chi$ be two constants. Let $t = v_1v_2v_3$ be a triangle with vertices on Σ . Assume that $d_{v_i}(v_i, z) \leq c_\omega\gamma$ and $d_{v_i}(v_i, z) \leq \rho \cdot \text{length}(t)$ for some point $z \in V_t(\{v_1, v_2, v_3\}) \cap \Sigma$. Let B_i be the ball centered at $\varphi_{v_i}(z)$ in the φ_{v_i} -space with radius $3 \cdot \text{length}(t)$. Then the following hold for sufficiently small ω .

- (i) For all point $y \in V_t(\{v_1, v_2, v_3\}) \cap \varphi_{v_i}^{-1}(B_i)$, the acute angle between $\mathbf{n}_{v_i}(t)$ and the tangent to $\varphi_{v_i}(V_t(\{v_1, v_2, v_3\}))$ at $\varphi_{v_i}(y)$ is $O(\omega)$.
- (ii) Define the function $\psi(c) = 2\tan(\sin^{-1}(c/2) + \beta(6\rho_1 c) + \alpha(6c))$. Let H be the plane of t . $V_t(\{v_1, v_2, v_3\}) \cap \varphi_{v_i}^{-1}(B_i)$ contains a curve segment η that contains z in its interior and intersects H exactly once at an endpoint. For any point $y \in \eta$, $d_{v_i}(y, z) \leq \psi(c_\omega) \cdot d_{v_i}(v_i, z)$.

Using Lemma 4.3, we can orient $\varphi_{v_3}(t_1)$ and $\varphi_{v_3}(t_2)$ in anti-clockwise order, when viewed from infinity in direction $\mathbf{n}_{v_3}(v_3)$, such that $\varphi_{v_3}(v_3v_4)$ receive opposite orientations from $\varphi_{v_3}(t_1)$ and $\varphi_{v_3}(t_2)$. This allows us to argue that $\varphi_{v_3}(t_1)$ and $\varphi_{v_3}(t_2)$ lie on opposite sides of a plane $\varphi_{v_3}(G)$ through $\varphi_{v_3}(v_3v_4)$ and parallel to $\mathbf{n}_{v_3}(v_3)$. Then the dihedral angle bound for $\varphi_{v_3}(t_1)$ and $\varphi_{v_3}(t_2)$ follows by applying Lemma 4.2(i). When we go back to the primal space, t_1 and t_2 also lie on opposite sides of G and the dihedral angle bound between t_1 and t_2 follows by Lemma 4.2(ii). The exact statements are given in the next lemma. Its proof is omitted.

LEMMA 4.4. *Let $\rho \in (1, \sqrt{2}]$ and $\rho_1 = 2/\chi$ be two constants. Let P be a set of vertices on Σ . Let $t_1 = v_1v_3v_4$ and $t_2 = v_2v_3v_4$ be two adjacent triangles where $v_i \in P$. Assume that for $1 \leq j \leq 2$, $V_{t_j}(P) \cap \Sigma$ is a single point z_j , $V_{v_3v_4}(P) \cap \Sigma$ contains only one curve joining z_1 and z_2 , and for any vertex v of t_j , $d_v(v, z_j) \leq c_\omega \gamma$ and $d_v(v, z_j) \leq \rho \cdot \text{length}(t_j)$. Then for sufficiently small ω , the dihedral angle between $\varphi_{v_3}(t_1)$ and $\varphi_{v_3}(t_2)$ as well as the dihedral angle between t_1 and t_2 are at least $\pi - 2\beta(4\rho_1 c_\omega) - 2\alpha(4c_\omega)$.*

5 Topological results

Consider the function that maps a point away from the medial axis of Σ to its nearest point on Σ . Let μ be the restriction of this function to $\text{Mesh } P$. The main results in this section are given in Theorem 5.1: μ is a homeomorphism between $\text{Mesh } P$ and Σ under suitable conditions. We will need a lower bound on the interpoint distances. Define

$$\epsilon_\omega = 1000\tau_\omega^2\rho_1^2\omega c_\omega.$$

The lower bound on interpoint distances is $\epsilon_\omega \gamma$. We first show in Lemma 5.1 below that μ is injective on the vertices in P .

LEMMA 5.1. *Let $\rho \in (1, \sqrt{2}]$ be a constant. Let P be a set of points on Σ . There exists a value ω_0 such that for any point $v \in P$, $\mu^{-1}(v) = \{v\}$, provided that the following conditions hold for some $\omega \leq \omega_0$.*

- For any points $a, b \in P$, $d_a(a, b) \geq \epsilon_\omega \gamma$.
- For any triangle $t = v_1v_2v_3$ in $\text{Mesh } P$, $d_{v_i}(v_i, z) \leq \rho \cdot \text{length}(t)$ and $d_{v_i}(v_i, z) \leq c_\omega \gamma$ for some point $z \in V_t(P) \cap \Sigma$.

Proof. (Sketch) Let m be the center of a medial ball M in the primal space that touches Σ at v . We go to the φ_v -space. Note that $\varphi_v(vm)$ lies on the normal line through $\varphi_v(v)$. Take a medial ball M' of $\varphi_v(\Sigma)$ at $\varphi_v(v)$ that intersects $\varphi_v(vm)$. Let $\varphi_v(m')$ be the center

of M' . Since the scaling factors satisfy the relation $k_{v,1} \leq k_{v,2} \leq k_{v,3}$, when we go back to the primal space, $\varphi_v^{-1}(M')$ is an empty ellipsoid centered at m' and it contains a ball centered at m' with radius $\|m' - v\|$. We conclude that $\|m' - v\| \leq \|m - v\|$, i.e., m' lies on mv .

It suffices to show that no triangle in $\text{Mesh } P$ intersects vm . Assume to the contrary that a triangle $t = v_1v_2v_3$ intersects vm . Let x be the point $t \cap vm$.

We argue that $\tau_{vv_i} \leq \tau_\omega$ for any vertex v_i of t . Starting with $d_{v_i}(v_i, v_j) \leq d_{v_i}(v_i, z) + d_{v_i}(v_j, z)$, we get

$$(5.1) \quad d_{v_i}(v_i, v_j) \leq d_{v_i}(v_i, z) + \tau_\omega^2 d_{v_j}(v_j, z) < 3c_\omega \gamma.$$

Since $x \in t$, there exists j such that $d_{v_i}(v_i, x) \leq d_{v_i}(v_i, v_j) < 3c_\omega \gamma$. So $x \in \Sigma^*$ and $\tau_{xv_i} \leq \tau_\omega$ by the definition of c_ω . By definition, $F_v = F_x$ and so $\tau_{vv_i} = \tau_{xv_i} \leq \tau_\omega$.

We claim that x lies between m' and v . Otherwise, in the primal space, since the distance from x to Σ is at least the distance from x to the boundary of the medial ball M centered at m , this distance is at least $\|m' - v\| = d_v(m', v)/\sqrt{k_{v,3}} \geq \gamma/\sqrt{k_{v,3}}$. But if we apply Lemma 4.2(iii) using any vertex of t , we can conclude that $\|x - v\| = O(c_\omega \gamma)$ which is less than $\gamma/\sqrt{k_{v,3}}$ for sufficiently small ω , a contradiction. This proves our claim. Hence, $\varphi_v(x)$ lies between $\varphi_v(m')$ and $\varphi_v(v)$.

We go back to the φ_v -space. Since M' is the medial ball of $\varphi_v(\Sigma)$, M' does not contain any vertex of $\varphi_v(t)$. By Lemma 4.3(ii), $V_t(\{v_1, v_2, v_3\})$ intersects the plane of t at a point q such that

$$(5.2) \quad d_{v_i}(v_i, q) \leq (1 + \psi(c_\omega))d_{v_i}(v_i, z) < 2c_\omega \gamma.$$

Take the smallest enclosing ball B of $\varphi_v(t)$ centered at $\varphi_v(q)$ in the φ_v -space. So the radius of this ball is $\max_{1 \leq i \leq 3} d_v(v_i, q) \leq \tau_\omega \cdot \max_{1 \leq i \leq 3} d_{v_i}(v_i, q) < 2\tau_\omega c_\omega \gamma$.

First, we consider the case in which B does not contain $\varphi_v(v)$. Since the diameter of B is no more than $4\tau_\omega c_\omega \gamma < \gamma$, B does not intersect the medial axis of $\varphi_v(\Sigma)$ and so $\varphi_v(m')$ lies outside B' . Thus the segment $\varphi_v(m'v)$ and $\varphi_v(t)$ lie on opposite sides of the bisector plane of B and M' . But then $\varphi_v(t)$ cannot intersect $\varphi_v(m'v)$ at $\varphi_v(x)$, a contradiction.

Second, we consider the case in which B contains $\varphi_v(v)$. So $d_v(v, v_i) \leq 4\tau_\omega c_\omega \gamma$. In the rest of the proof, we show that this leads to the contradiction that $d_v(v, z) < d_{v_j}(v_j, z)$ for some j . Lemma 4.2(iii) implies that $d_v(x, v) < 640\tau_\omega^2\rho_1^2c_\omega^2\gamma$. For $1 \leq i \leq 3$, since $d_v(v, v_i) \geq \epsilon_\omega \gamma$, we get $d_v(x, v) \leq 640\tau_\omega^2\rho_1^2 \cdot (c_\omega^2/\epsilon_\omega) \cdot d_v(v, v_j)$. By definition, $\epsilon_\omega = 1000\tau_\omega^2\rho_1^2\omega c_\omega$ and $c_\omega \leq \omega^3$, so we obtain

$$(5.3) \quad d_v(x, v) < \omega^2 \cdot d_v(v, v_i).$$

Next, we bound the distance $d_v(x, q)$. First, for any v_i ,

$$d_v(v_i, x) \geq d_v(v, v_i) - d_v(v, x) \geq (1 - \omega^2) \cdot d_v(v, v_i).$$

Observe that, as $\varphi_v(x)$ lies inside $\varphi_v(t)$, there is always a vertex v_j such that $\varphi_v(\angle q x v_j) \geq \pi/2$. Therefore,

$$\begin{aligned} d_v(x, q) &\leq \sqrt{d_v(v_j, q)^2 - d_v(v_j, x)^2} \\ &\leq \sqrt{\tau_\omega^2 d_{v_j}(v_j, q)^2 - (1 - \omega^2)^2 d_v(v, v_j)^2}. \end{aligned}$$

Since $d_{v_j}(v_j, q) \leq 2c_\omega\gamma$ by equation (5.2) and $d_v(v, v_j) \geq \epsilon_\omega\gamma$ by assumption, we have $d_v(v, v_j) \geq \epsilon_\omega/(2c_\omega) \cdot d_{v_j}(v_j, q)$. Using λ to denote $1000\tau_\omega^2\rho_1^2$, we get $d_v(v, v_j) \geq (\lambda\omega/2) \cdot d_{v_j}(v_j, q) > \tau_\omega\lambda\omega/(2\sqrt{2}) \cdot d_{v_j}(v_j, q)$. Thus,

$$d_v(x, q) \leq \tau_\omega d_{v_j}(v_j, q) \cdot \sqrt{1 - (1 - \omega^2)^2 \cdot (\lambda^2\omega^2/8)}$$

For sufficiently small ω , this simplifies to

$$d_v(x, q) \leq \tau_\omega(1 - \lambda^2\omega^2/32) \cdot d_{v_j}(v_j, q).$$

Since $\lambda = 1000\tau_\omega^2\rho_1^2$, $\lambda^2\omega^2/64 \geq 4\omega^3$. Also, since $\psi(c_\omega) = O(c_\omega) = O(\omega^3)$, $\lambda^2\omega^2/64 \geq 2\psi(c_\omega)$ for sufficiently small ω . Hence, we obtain

$$d_v(x, q) \leq \tau_\omega(1 - 4\omega^2 - 2\psi(c_\omega)) \cdot d_{v_j}(v_j, q).$$

Recall that the radius of B is equal to $d_v(v_k, q)$ for some v_k . By equation (5.3), $d_v(v, x) \leq \omega^2 \cdot d_v(v, v_i)$ for any vertex v_i . Thus $d_v(v, x) \leq \omega^2 \cdot (d_v(v, q) + d_v(v_i, q)) \leq 2\omega^2 \cdot \text{radius}(B) = 2\omega^2 \cdot d_v(v_k, q) \leq 2\tau_\omega\omega^2 \cdot d_{v_k}(v_k, q) = 2\tau_\omega\omega^2 \cdot d_{v_j}(v_j, q)$. It follows that

$$\begin{aligned} d_v(v, q) &\leq d_v(v, x) + d_v(x, q) \\ &\leq \tau_\omega(1 - 2\omega^2 - 2\psi(c_\omega)) \cdot d_{v_j}(v_j, q). \end{aligned}$$

By Lemma 4.3(ii), $d_{v_j}(q, z) \leq \psi(c_\omega) \cdot d_{v_j}(v_j, z)$. So $d_v(q, z) \leq \tau_\omega\psi(c_\omega) \cdot d_{v_j}(v_j, z)$. Hence,

$$\begin{aligned} d_v(v, z) &\leq d_v(v, q) + d_v(q, z) \\ &< \tau_\omega(1 - 2\omega^2) \cdot d_{v_j}(v_j, z). \end{aligned}$$

Since $\tau_\omega = 1/(1 - \omega^2)$, $\tau_\omega(1 - 2\omega^2) < 1$. So $d_v(v, z) < d_{v_j}(v_j, z)$ which contradicts the fact that $z \in V_t(P)$. $\square$

In the next result, when we say Mesh P is a 2-manifold combinatorially, we mean that the incidence relations among the triangles indicate a 2-manifold, but we do not care whether Mesh P self-intersects.

THEOREM 5.1. *Let $\rho \in (1, \sqrt{2}]$ be a constant. Let P be a set of points on Σ . There exists a value ω_0 such that Mesh P does not self-intersect and Mesh P is homeomorphic to $\mu(\text{Mesh } P)$, provided that the following conditions hold for some $\omega \leq \omega_0$.*

- Mesh P is combinatorially a 2-manifold.

- For each triangle t in Mesh P , $V_t(P) \cap \Sigma$ is a single point. For each edge e in Mesh P , $V_e(P) \cap \Sigma$ is a simple open curve.
- For any points $a, b \in P$, $d_a(a, b) \geq \epsilon_\omega\gamma$. For any triangle $t = v_1v_2v_3$ in Mesh P , $d_{v_i}(v_i, z) \leq \rho \cdot \text{length}(t)$ and $d_{v_i}(v_i, z) \leq c_\omega\gamma$, where $z = V_t(P) \cap \Sigma$.

Hence, under the above conditions, if Mesh P contains a vertex in each connected component of Σ , then Mesh P is homeomorphic to Σ .

Proof. (Sketch) Let ω_0 be a small enough value for which the previous results hold. By our assumption, it can be checked that Mesh P avoids the medial axis of Σ . We show that μ is a homeomorphism between Mesh P and $\mu(\text{Mesh } P)$. Since Mesh P is compact and μ is continuous, $\mu(\text{Mesh } P)$ is also compact. Thus it suffices to show that μ is injective.

First, we show that μ is injective on each triangle t in Mesh P . Assume to the contrary that there is a point $x \in \Sigma$ such that μ maps two distinct points p and q in a triangle t to x . So p, q and x lie on the line normal to Σ at x . Let v be a vertex of t . Let z be the point $V_t(P) \cap \Sigma$. For any other vertex w of t , $d_v(v, w) \leq d_v(v, z) + \tau_*d_w(w, z) = O(c_\omega\gamma)$. By Lemma 4.2(iii), the distance from p to Σ is $O(c_\omega\gamma)$. Thus $\|p - x\| = O(c_\omega\gamma)$. So for sufficiently small ω , $p \in \Sigma^*$ and so F_p is defined and $F_p = F_x$ by definition. Since $\|p - x\| = O(c_\omega\gamma)$, $d_p(p, x) = d_x(p, x) = \sqrt{k_{x,3}} \cdot \|p - x\| = O(\sqrt{k_{x,3}} \cdot c_\omega\gamma) = O(\sqrt{c_\omega\gamma})$ for sufficiently small ω . Since $p \in t$, there is a vertex w of t such that $d_v(v, p) \leq d_v(v, w)$ which has been shown to be $O(c_\omega\gamma)$. We conclude that $d_v(v, x) \leq d_v(v, p) + d_v(p, x) \leq O(c_\omega\gamma) + \tau_*d_p(p, x) = O(\sqrt{c_\omega\gamma})$. By Lemma 3.2(ii), $\mathbf{n}(v)$ and $\mathbf{n}(x)$ are nearly parallel. Then by Lemma 4.2(ii), t and $\mathbf{n}(x)$ are nearly orthogonal. But since pq lies on t , t must be parallel to $\mathbf{n}(x)$, a contradiction.

Second, we show that μ is injective. For each point $x \in \mu(\text{Mesh } P)$, define $h(x)$ to be the number of points mapped to x by μ . One can visualize $h(x)$ as the number of intersection points between Mesh P and the line segment s_x joining the centers of the two medial balls at x . Since μ is injective on each triangle, $h(x)$ is a finite number and as x varies over Σ , $h(x)$ stays constant if s_x does not cross any triangle edge. Suppose that $h(x)$ changes when s_x crosses a triangle edge. Since Mesh P is combinatorially a 2-manifold, this edge is incident to two triangles t_1 and t_2 . The change in $h(x)$ implies that $\mu(t_1)$ and $\mu(t_2)$ overlap. But this contradicts the large dihedral angle between t_1 and t_2 (Lemma 4.4) and the fact that t_1 and t_2 are almost perpendicular to the normal of any common vertex (Lemma 4.2(ii)). Thus h is a constant function.

Then Lemma 5.1 implies that $h = 1$. It follows that μ is a homeomorphism. $\square$

6 Anisotropic meshing algorithm

We first discuss some primitives to be used by our algorithm ANISOMESH. Given a sample set P , we assume an incremental algorithm to compute $\text{Vor } P$, e.g. the incremental lower envelope algorithm in [9]. The lower envelope can be used to answer which Voronoi edge, facet, or cell contains a point x in $\mathbb{R}^3$. The *critical points* of a curve or surface with respect to a direction d are the points at which d is normal to the curve or surface. We use $E : \mathbb{R}^3 \rightarrow \mathbb{R}$ to denote the implicit function for Σ , i.e., $x \in \Sigma$ iff $E(x) = 0$. For any $v, w \in P$, let $H_{vw}(x) = 0$ denote the equation of $V_{vw}(\{v, w\})$. We assume a numerical solver for a system of equations. For example, we can then compute the intersection points between Σ and a Voronoi edge of $\text{Vor } P$. ANISOMESH uses the following three routines. Let x_1, x_2 , and x_3 be the coordinate axes. Given any equation $G(x) = 0$, by taking partial derivatives, we get $G'(x) = (\frac{\partial G(x)}{\partial x_1}, \frac{\partial G(x)}{\partial x_2}, \frac{\partial G(x)}{\partial x_3})$, which is the equation of a normal at x to $G(x)$.

- **FacetTest**(edge v_1v_2 in $\text{Mesh } P$): Compute the vector $d = \mathbf{n}(v_1) \times (v_2 - v_1)$. If $V_{v_1v_2}(P) \cap \Sigma$ contains any critical point with respect to d or if $V_{v_1v_2}(P)$ is tangent to Σ at some point, return one as follows. The intersection $V_{v_1v_2}(\{v_1, v_2\}) \cap \Sigma$ is a collection of disjoint smooth closed curves. The equation of the tangent at any point $x \in V_{v_1v_2}(\{v_1, v_2\}) \cap \Sigma$ is $H'_{v_1v_2}(x) \times E'(x)$. The set of critical points on $V_{v_1v_2}(\{v_1, v_2\}) \cap \Sigma$ with respect to d is defined by: $E(x) = 0, H_{v_1v_2}(x) = 0, \langle H'_{v_1v_2}(x) \times E'(x), d \rangle = 0$. Solve it and return any solution that lies on $V_{v_1v_2}(P)$. The tangency case is handled similarly. At the contact point x , $H'_{v_1v_2}(x) \times E'(x) = 0$. Thus, solve the system: $E(x) = 0, H_{v_1v_2}(x) = 0, H'_{v_1v_2}(x) \times E'(x) = 0$, and return any solution that lies on $V_{v_1v_2}(P)$.
- **PatchMax**(vertex v in $\text{Mesh } P$): Return the point x in $V_v(P) \cap \Sigma$ that maximizes $d_v(v, x)$ as follows. We first handle points on the boundary of $V_v(P) \cap \Sigma$. The intersection points on the edges of $V_v(P)$ can be computed directly. Take a Voronoi neighbor w of v . If the maximum is achieved at some interior point $x \in V_{vw}(P) \cap \Sigma$, the tangent to $\varphi_v(V_{vw}(P) \cap \Sigma)$ at $\varphi_v(x)$ is orthogonal to the vector $\varphi_v(x - v)$. The equation of the tangent to $\varphi_v(V_{vw}(P) \cap \Sigma)$ at $\varphi_v(x)$ is $F_v \cdot (H'_{vw}(x) \times E'(x))$. Solve the system: $E(x) = 0, H_{vw}(x) = 0, \langle F_v \cdot (x - v), F_v \cdot (H'_{vw}(x) \times E'(x)) \rangle = 0$. Collect the solutions that lie on

$V_{vw}(P)$ and repeat for other Voronoi neighbors of v . If the maximum is achieved at some interior point $x \in V_v(P) \cap \Sigma$, $\mathbf{n}_v(x)$ makes an angle 0 or π with the vector $\varphi_v(x - v)$. The equation of the normal to $\varphi_v(\Sigma)$ at $\varphi_v(x)$ is $F_v^{-1} \cdot E'(x)$. Solve the system: $E(x) = 0, (F_v \cdot (x - v)) \times (F_v^{-1} \cdot E'(x)) = 0$. Collect the solutions that lie inside $V_v(P)$. After collecting all the candidate points, return the point x that maximizes $d_v(v, x)$.

- **SilhTest**(vertex v in $\text{Mesh } P$): If the set $\{x \in V_v(P) \cap \Sigma : \angle \mathbf{n}(x), \mathbf{n}(v) = \pi/2\}$ is non-empty, return a point in it as follows. The larger set $J = \{x \in \Sigma : \angle \mathbf{n}(x), \mathbf{n}(v) = \pi/2\}$ is a collection of disjoint smooth closed curves [14]. The set J satisfies the system: $E(x) = 0, K(x) = 0$, where $K(x) = \langle E'(x), \mathbf{n}(v) \rangle$. We check whether J intersects the boundary of $V_v(P)$ by solving the system: $E(x) = 0, K(x) = 0, H_{vw}(x) = 0$, for all Voronoi neighbors w of v . If there is a solution lying on $V_{v,w}(P)$, return it. Otherwise, let d be any direction orthogonal to $\mathbf{n}(v)$. If some component of J lies completely inside $V_v(P)$, J has a critical point with respect to d inside $V_v(P)$. The tangent to J at x is given by $K'(x) \times E'(x)$. Solve the system: $E(x) = 0, K(x) = 0, \langle K'(x) \times E'(x), d \rangle = 0$. If there is a solution lying inside $V_v(P)$, return it.

The SilhTest routine was first introduced in [11] for isotropic surface meshing. We are ready to describe our algorithm to mesh Σ .

ANISOMESH(Σ):

1. Initialize P to contain one critical point of Σ with respect to the vertical direction. (Even though Σ may have multiple components, one critical point is enough for initialization.)
2. Repeatedly apply a rule from the following list to insert a new point into P until no rule is applicable. A rule with smaller index has higher priority. Let $\rho \in (\tau_*, \sqrt{2}]$, $\zeta \in (0, \pi/4)$ and $\ell > 0$ be some constants.
 - (a) If there is a triangle $t = v_1v_2v_3$ in $\text{Mesh } P$ such that $d_{v_i}(v_i, x) > \rho \cdot \text{length}(t)$ for some point $x \in V_t(P) \cap \Sigma$, insert x .
 - (b) If a Voronoi edge $V_{v_1v_2v_3}(P)$ intersects Σ at least twice or tangentially, insert the point $x \in V_{v_1v_2v_3}(P) \cap \Sigma$ that maximizes $d_{v_i}(v_i, x)$.
 - (c) If **FacetTest**(v_1v_2) returns a point x for some edge v_1v_2 in $\text{Mesh } P$, insert x .
 - (d) If the incident triangles of a vertex v in $\text{Mesh } P$ do not form a disk combinatorially, insert the point x

returned by PatchMax(v). (By not being a disk combinatorially, we mean the incidence relations among the incident triangles do not indicate a topological disk, and we do not care whether some triangles cross each other.)

- (e) If Mesh P self-intersects, find the triangle $t = v_1 v_2 v_3$ in Mesh P such that $d_{v_i}(v_i, x)$ is maximum, where $x = V_t(P) \cap \Sigma$, and insert x .
- (f) If SilhTest(v) returns a point x for some vertex v of Mesh P , insert x .
- (g) If there is a triangle $t = v_1 v_2 v_3$ in Mesh P such that $\angle \mathbf{n}_{v_i}(t), \mathbf{n}_{v_i}(v_i)$ or $\angle \mathbf{n}(t), \mathbf{n}(v_i)$ exceeds ζ for some $1 \leq i \leq 3$, insert the point $V_t(P) \cap \Sigma$.
- (h) If $d_{v_1}(v_1, v_2) > \ell$ for some edge $v_1 v_2$ in Mesh P , take a triangle t incident to $v_1 v_2$ and insert the point $V_t(P) \cap \Sigma$.

Except for the upper bound on triangle size, all conditions in Theorem 5.1 are enforced in steps 2(a)–(e). Step 2(c) is a sufficient condition to ensure that each Voronoi facet does not intersect in a closed curve. Then step 2(d) ensures that each Voronoi facet intersects Σ in a simple open curve, and the boundary of $V_v(P) \cap \Sigma$ is a single closed curve for each $v \in P$. After step 2(e), tunnels or isolated components in $V_v(P) \cap \Sigma$ may still exist and step 2(f) detects that. Previous experience with the isotropic surface meshing problem [11] suggests that even if the triangulation is homeomorphic to Σ , it may not be a good geometric approximation of Σ . Therefore, steps 2(g) and 2(h) are introduced to enforce user-supplied upper bounds on the maximum angle deviation and maximum edge length. The termination of ANISOMESH follows from establishing a lower bound on the interpoint distances throughout the algorithm.

LEMMA 6.1. *Let $\omega < 1/(1000\tau_*^3\rho_1^2)$ be a small enough constant such that Theorem 5.1 holds, $\beta(6\rho_1\tau_*\epsilon_\omega) + \alpha(6\tau_*\epsilon_\omega) < \zeta$, and $3\tau_*\epsilon_\omega\gamma < \ell$. After each point insertion in ANISOMESH, $d_a(a, b) \geq \epsilon_\omega\gamma$ for any vertices $a, b \in P$.*

Proof. (Sketch) The lemma is true after step (1). We prove the lemma inductively. It suffices to show that $d_v(v, x) \geq \tau_*\epsilon_\omega\gamma$ if a point $x \in V_v(P) \cap \Sigma$ is inserted. Then $d_x(w, x) \geq \epsilon_\omega\gamma$ for all $w \in P$; otherwise, $d_w(w, x) \leq \tau_*d_x(w, x) < \tau_*\epsilon_\omega\gamma \leq d_v(v, x)$, a contradiction.

If step 2(a) inserts a point $x \in V_t(P) \cap \Sigma$, then for any vertex v_i of t , $d_{v_i}(v_i, x) > \rho \cdot \text{length}(t) \geq \rho\epsilon_\omega\gamma > \tau_*\epsilon_\omega\gamma$ as $\rho > \tau_*$.

Suppose that step 2(b) inserts a point in $V_t(P) \cap \Sigma$, where $t = v_1 v_2 v_3$. Assume to the contrary that

$d_{v_i}(v_i, z) < \tau_*\epsilon_\omega\gamma$ for all points $z \in V_t(P) \cap \Sigma$. Suppose that $V_t(P)$ intersects Σ at two points z_1 and z_2 . The case where $V_t(P)$ meets Σ tangentially is easier. We go to the φ_{v_1} -space. Let B_1 be the ball centered at $\varphi_{v_1}(z_1)$ in the φ_{v_1} -space with radius $3 \cdot \text{length}(t) \geq 3\epsilon_\omega\gamma$. Let B and B' be the two medial balls of $\varphi_{v_1}(\Sigma)$ meeting at $\varphi_{v_1}(z_1)$. By Lemma 3.2(ii) and Lemma 4.2(i), $\mathbf{n}_v(z_1)$ and $\mathbf{n}_v(t)$ are nearly parallel. Then by Lemma 4.3(i), $\varphi_v(V_t(P)) \cap B_1$ lies close to the normal at $\varphi_{v_1}(z_1)$. So $\varphi_v(V_t(P)) \cap B_1$ lies inside $B \cup B'$, which implies that $d_{v_1}(z_1, z_2) \geq \text{radius}(B_1) \geq 3\epsilon_\omega\gamma$, contradicting $d_{v_1}(z_1, z_2) \leq d_{v_1}(v_1, z_1) + d_{v_1}(v_1, z_2) < 2\tau_*\epsilon_\omega\gamma$.

Suppose that FacetTest($v_1 v_2$) returns a point x in step 2(c). We skip the tangency case as the analysis is similar. So some plane H orthogonal to $\mathbf{n}(v_1) \times (v_2 - v_1)$ meets $V_{v_1 v_2}(P) \cap \Sigma$ tangentially at x . Then $\varphi_{v_1}(H)$ is orthogonal to $\mathbf{n}_{v_1}(v_1) \times \varphi_{v_1}(v_2 - v_1)$ and it meets $\varphi_{v_1}(V_{v_1 v_2}(P) \cap \Sigma)$ tangentially at $\varphi_{v_1}(x)$. Assume to the contrary that $d_{v_1}(v_1, x) < \tau_*\epsilon_\omega\gamma$. We go to the φ_{v_1} -space. It can be checked that $d_{v_1}(v_1, v_2) < 3\tau_*\epsilon_\omega\gamma$. Let e_x be a normal to $\varphi_{v_1}(V_{v_1 v_2}(P))$ at $\varphi_{v_1}(x)$ pointing towards the side of $\varphi_{v_1}(v_2)$. Using an analysis in proving Lemma 4.3(i), one can show that e_x is nearly parallel to $\varphi_{v_1}(v_2 - v_1)$. By Lemma 3.2(ii), $\mathbf{n}_{v_1}(x)$ is nearly parallel to $\mathbf{n}_{v_1}(v_1)$. By Lemma 3.2(i), the vector $\varphi_{v_1}(v_2 - v_1)$ is nearly orthogonal to $\mathbf{n}_{v_1}(v_1)$. So $\mathbf{n}_{v_1}(x) \times e_x$ is nearly parallel to $\mathbf{n}_{v_1}(v_1) \times \varphi_{v_1}(v_2 - v_1)$, which is normal to $\varphi_{v_1}(H)$. But since $\mathbf{n}_{v_1}(x) \times e_x$ is tangent to $\varphi_{v_1}(V_{v_1 v_2}(P) \cap \Sigma)$ at $\varphi_{v_1}(x)$, $\mathbf{n}_{v_1}(x) \times e_x$ is parallel to $\varphi_{v_1}(H)$, a contradiction.

We omit the analysis of step 2(d). Suppose that step 2(e) inserts the point $x = V_t(P) \cap \Sigma$ for some triangle $t = v_1 v_2 v_3$. Before the insertion of x , Mesh P satisfies all the conditions in Theorem 5.1 except the upper bound on triangle size. Thus $d_{v_i}(v_i, x) > c_\omega\gamma = \epsilon_\omega\gamma/(1000\tau_*^2\rho_1^2\omega) > \tau_*\epsilon_\omega\gamma$.

Suppose that step 2(f) inserts the point $x \in V_v(P)$. So $\angle \mathbf{n}(v), \mathbf{n}(x) = \pi/2$ which implies that $\angle \mathbf{n}_v(v), \mathbf{n}_v(x) = \pi/2$. Thus $d_v(v, x)$ must be greater than $\tau_*\epsilon_\omega\gamma$ by Lemma 3.2(ii).

Suppose that step 2(g) inserts the point $x = V_t(P) \cap \Sigma$ for $t = v_1 v_2 v_3$. So $\angle \mathbf{n}_{v_i}(t), \mathbf{n}_{v_i}(v_i)$ or $\angle \mathbf{n}(t), \mathbf{n}(v_i)$ exceeds ζ for some i . If $d_{v_i}(v_i, x) < \tau_*\epsilon_\omega\gamma$, it can be checked that $d_{v_i}(v_i, v_j) < 3\tau_*\epsilon_\omega\gamma$. But then Lemma 4.2(i) and (ii) imply that both $\angle \mathbf{n}_{v_i}(t), \mathbf{n}_{v_i}(v_i)$ and $\angle \mathbf{n}(t), \mathbf{n}(v_i)$ are no more than $\beta(6\rho_1\tau_*\epsilon_\omega) + \alpha(6\tau_*\epsilon_\omega) < \zeta$, a contradiction.

Suppose that step 2(h) finds an edge vw such that $d_v(v, w) > \ell$. Let z be the point $V_t(P) \cap \Sigma$ for some triangle t incident to vw . Then $d_v(v, z) \geq \tau_*\epsilon_\omega\gamma$; otherwise, $d_v(v, w) \leq d_v(v, z) + d_v(z, w) \leq d_v(v, z) + \tau_*^2 d_w(w, z) < 3\tau_*\epsilon_\omega\gamma < \ell$, a contradiction. $\square$

THEOREM 6.1. *Let $\rho \in (\tau_*, \sqrt{2}]$, $\ell > 0$, $\zeta \in (0, \pi/4)$, and $\chi = \frac{1-32(\tau_*^4-1)}{128\tau_*^2}$ be some constants. Assume that Σ is a smooth closed implicit surface and there is an oracle to return the 2D metric tensor at any point $x \in \Sigma$. ANISOMESH(Σ) samples a set P of vertices on Σ and computes a triangulation $\text{Mesh } P$ homeomorphic to Σ . For any triangle t in $\text{Mesh } P$ and for any edge vw of t , $d_v(v, w) \leq \ell$, $\angle(\mathbf{n}(t), \mathbf{n}(v)) \leq \zeta$, $\angle_{\mathbf{n}_v}(t), \mathbf{n}_v(v) \leq \zeta$, and each angle of $\varphi_v(t)$ is at least $\sin^{-1}(\chi)$.*

Proof. (Sketch) The geometric approximation is guaranteed by previous results. Let S_v denote $V_v(P) \cap \Sigma$. By the end of ANISOMESH, S_v is connected and ∂S_v is a single closed curve. Then it has been shown that S_v is a topological disk and it projects to a topological disk on the tangent plane at v [11]. We refine ∂S_v by inserting a vertex in the interior of each boundary edge $V_{vw}(P) \cap \Sigma$ of S_v . Then we refine $\text{Mesh } P$ into a triangulation T by refining each triangle using a barycentric subdivision. Thanks to the barycentric subdivision, for each vertex $v \in P$, there is a bijection between the edges in its link in T (i.e., the boundary of the fan of triangles incident to v) and the refined ∂S_v . Afterwards, this boundary homeomorphism can be extended to a homeomorphism between the fan of triangles incident to v in T and S_v . Stitching these homeomorphisms together yields a homeomorphism between Σ and $\text{Mesh } P$. $\square$

7 Conclusion

There are several directions for future work. We plan to conduct an experimental study of the effectiveness of the algorithm. There has been some recent work on remeshing a given surface mesh in the anisotropic setting, e.g. [1], and this problem awaits a thorough theoretical study. This has been done in the isotropic case by Dey, Li, and Ray [13]. The problem of generating a 3D anisotropic volume mesh remains open.

References

- [1] P. Alliez, D. Cohen-Steiner, O. Devillers, B. Levy, and M. Desbrun. Anisotropic polygonal remeshing. *ACM Transactions on Graphics*, 22 (2003), 485–493.
- [2] N. Amenta and M. Bern. Surface reconstruction by Voronoi filtering. *Discr. Comput. Geom.*, 22 (1999), 481–504.
- [3] N. Amenta, S. Choi, T. K. Dey and N. Leekha. A simple algorithm for homeomorphic surface reconstruction. *Internat. J. Comput. Geom. Applications*, 12 (2002), 125–141.
- [4] J. Bloomenthal. Polygonization of implicit surfaces. *Proc. Comput. Aided Geom. Design*, 5 (1988), 341–355.
- [5] J.-D. Boissonnat and F. Cazals. Natural neighbor coordinates of points on a surface. *Comput. Geom. Theory Appl.*, 19 (2001), 87–120.
- [6] J.-D. Boissonnat and S. Oudot. Provably good surface sampling and approximation. *Eurographics Sympos. Geom. Process.* (2003), 9–18.
- [7] J.-D. Boissonnat, D. Cohen-Steiner and G. Vegter. Isotopic implicit surface meshing. *Proc. 36th ACM Sympos. Theory Comput.*, 2004, 301–309.
- [8] J.-D. Boissonnat, C. Wormser, and M. Yvinec. Anisotropic diagrams: Labelle Shewchuk approach revisited. *CCCG* 2005.
- [9] B. Chazelle, H. Edelsbrunner, L. Guibas, and M. Sharir. A singlyexponential stratification scheme for real semi-algebraic varieties and its applications. *Theoret. Comput. Sci.*, 84 (1991), 77–105.
- [10] H.-L. Cheng, T. K. Dey, H. Edelsbrunner and J. Sullivan. Dynamic skin triangulation. *Discrete Comput. Geom.*, 25 (2001), 525–568.
- [11] S.-W. Cheng, T. K. Dey, E. A. Ramos, and T. Ray. Sampling and meshing a surface with guaranteed topology and geometry. *Proc. 20th Annu. Sympos. Comput. Geom.*, 2004, 280–289.
- [12] J.C. Cuilliere. An adaptive method for the automatic triangulation of 3D parametric surfaces. *Computer Aided Design*, 30 (1998), 139–149.
- [13] T.K. Dey, G. Li, and T. Ray. Polygonal surface remeshing with Delaunay refinement. *Proc. 14th International Meshing Roundtable*, 2005, 343–351.
- [14] H. Edelsbrunner and J. Harer. Jacobi sets of multiple Morse functions. In *Foundations of Computational Mathematics*, eds. F. Cucker, R. DeVore, P. Olver and E. Sueli, Cambridge Univ. Press, 2003, 37–57.
- [15] T. Karkanis and A.J. Stewart. Curvature-dependent triangulation of implicit surfaces. *IEEE Computer Graphics and Applications*, 21 (2001), 60–69.
- [16] F. Labelle and J.R. Shewchuk. Anisotropic Voronoi diagrams and guaranteed-quality anisotropic mesh generation. *Proc. 19th Annu. Sympos. Comput. Geom.*, 2003, 191–200.
- [17] T.S. Lau and S.H. Lo. Finite element mesh generation over analytical surfaces. *Computers and Structures*, 59 (1996), 301–309.
- [18] G. Leibon and D. Letscher. Delaunay triangulations and Voronoi diagrams for Riemannian manifolds. *Proc. 16th Annu. Sympos. Comput. Geom.*, 2000, 341–349.
- [19] W.E. Lorensen and H.E. Cline. Marching cubes: a high resolution 3D surface reconstruction algorithm. *Computer Graphics (SIGGRAPH '87 Proceedings)*, 21 (1987), 163–170.
- [20] M. Sharir. Almost tight bounds for lower envelopes in higher dimensions. *Discrete & Comput. Geom.*, 12 (1994), 327–345.
- [21] J.R. Tristano, S.J. Owen, and S.A. Canann. Advancing front surface mesh generation in parametric space using a Riemannian surface definition. *Proc. 7th Intl. Meshing Roundtable* 1998.